

# Dicyclic Groups and Frobenius Manifolds

Yassir Dinar and Zainab Al-Maamari

Department of Mathematics, College of Science, Sultan Qaboos University, P.O. Box 36, Al-Khod, PC 123, Muscat, Sultanate of Oman. \*Email: dinar@squ.edu.om

**ABSTRACT:** The orbits space of an irreducible representation of a finite group is a variety whose coordinate ring is finitely generated by homogeneous invariant polynomials. Boris Dubrovin showed that the orbits spaces of the reflection groups acquire the structure of polynomial Frobenius manifolds. Dubrovin's method to construct examples of Frobenius manifolds on orbits spaces was carried for other linear representations of discrete groups which have in common that the coordinate rings of the orbits spaces are polynomial rings. In this article, we show that the orbits space of an irreducible representation of a dicyclic group acquires two structures of Frobenius manifolds. The coordinate ring of this orbits space is not a polynomial ring.

**Keywords:** Differential geometry; Frobenius manifolds; Invariant theory; Dicyclic group; Orbifolds.

زمر مزدوجة الدورة وفضاءات فروبينيس الهندسية

ياسر دینار و زينب المعمری

**الملخص:** إن فضاء المدارات للتمثيل غير القابل للاختزال لزمرة محدودة هو فضاء متنوع يمتلك حلقة إحداثية محدودة تتشكل من كثيرات حدود متجانسة وغير متبدلة. أثبت بوريس دوبروفين أن الفضاءات المدارية للزمر المتعكسة تحتفظ ببنية فضاءات فروبينيس الهندسية لكثيرات الحدود. باستخدام طريقة دوبروفين، يمكن إيجاد أمثلة لفضاءات فروبينيس على فضاءات مدارية تمثلت خطياً لزمر متقطعة أخرى، لها خاصية عامة في الحلقات الإحداثية للفضاءات المدارية، تكون حلقات كثيرة الحدود. نبين في هذه المقالة، أن فضاء المدارات لتمثيلات غير قابلة للاختزال لزمر ثنائية الدوران تكتسب هيكلين من فضاءات فروبينيس. إن الحلقة الإحداثية لفضاء المدارات المذكور ليست حلقة لكثيرات الحدود.

**الكلمات المفتاحية:** الهندسة التفاضلية، فضاءات فروبينيس، النظرية الثابتة، مجموعة ثنائية الدوران، المدارات.



## 1. Introduction

The notion of a Frobenius manifold was introduced by Boris Dubrovin as a geometric realization of a potential  $F$  satisfying a system of partial differential equations known in topological fields theory as WDVV equations [1].

Besides topological fields theory, Frobenius manifolds appear in many fields such as invariant theory, integrable systems, quantum cohomology and singularity theory. This article contributes to the relation between Frobenius manifolds and invariant theory.

Let  $W$  be a finite group of linear transformations acting on a complex vector space  $V$  of dimension  $r$ . Then the orbits space  $M = V/W$  of this group is a variety whose coordinate ring is the ring of invariant polynomials  $C[V]^W$ . The ring  $C[V]^W$  is finitely generated by homogeneous polynomials. If  $f_1, f_2, \dots, f_m$  is a set of such generators then  $m \geq r$  and the relation between them is called syzygies. The set of generators are not unique, nor are their degrees [2,3].

An element  $w \in W$  is called a reflection if it fixes a subspace of  $V$  of codimension one pointwise.

The group  $W$  is called a complex reflection group if it is generated by reflections. Then Shephard-Todd-Chevalley theorem states that  $W$  is a reflection group if and only if the invariant ring  $C[V]^W$  is a polynomial ring [11], i.e. it is generated by  $r$  algebraically independent homogeneous polynomials (so there are no syzygies). Furthermore, when  $W$  is a reflection group, the degrees of such a set generators of  $C[V]^W$  are uniquely specified by the group and we refer to them as the degrees of  $W$ .

Let us assume  $W$  is a Shephard group, i.e. a symmetry group of a regular complex polytope. Then  $W$  is a reflection group. Let  $f_1, f_2, \dots, f_r$  be a set of algebraically independent homogeneous generators of  $C[V]^W$ . We assume that degree  $f_i$  is less than or equal to the degree  $f_j$  when  $i < j$ . Then the inverse of the Hessian of  $f_1$  defines a flat metric  $(\cdot, \cdot)_2$  on  $T^*M$  [4]. There is another flat metric  $(\cdot, \cdot)_1$  on  $T^*M$ , which was studied initially by Saito [5, 6], defined as the Lie derivative of  $(\cdot, \cdot)_2$  along the vector field  $e = \partial_{f_r}$ . The two metrics form what is called ‘a flat pencil of metrics’ (more details are given below). Dubrovin used the properties of this flat pencil of metrics to construct polynomial Frobenius manifolds [7] (see [8] and [9] for the case of Coxeter groups). This article is about applying Dubrovin’s method for other finite linear groups than Shephard groups.

Dubrovin’s method to construct Frobenius manifolds, through finding flat pencils of metrics on orbits spaces, was carried out for infinite linear groups like extended affine Weyl groups [10, 11], Jacobi groups [12] and recently a new extension of affine Weyl groups [13]. They all have in common that the invariant rings are polynomial rings. Moreover, even when considering a generalization of Frobenius manifold structure on orbits spaces, many results were obtained under the assumption that the invariant ring is a polynomial ring [14]. It is then a natural question to ask about applying Dubrovin’s method on orbits spaces of finite non-reflection groups.

In this article we apply Dubrovin’s method and construct Frobenius manifolds on orbits spaces of Dicyclic groups. The resulting Frobenius manifolds can be obtained by using an ad-hoc procedure, but it is fascinating to find them on orbits spaces of some group. Precisely, we will show that the orbits space of the Dicyclic group of order  $4n$  is endowed with two structures of Frobenius manifolds which up to scaling has the following potential

$$\mathbb{F}(z_1, z_2) = z_1^k + \frac{1}{2}z_2^2 z_1, \quad k = \frac{3-d}{1-d} \tag{1.1}$$

where  $d$  is  $\frac{2+\sqrt{3}n}{\sqrt{3}n}$  or  $\frac{2-\sqrt{3}n}{\sqrt{3}n}$ .

To make the article as self-contained as possible, we review in the next section the definition of the Frobenius manifold and its relation with the theory of flat pencils of metrics. In the last section we obtain the promised Frobenius manifolds by direct calculations.

## 2. Preliminaries

### 2.1 Frobenius manifolds

A Frobenius algebra is a commutative associative algebra with unity  $e$  and an invariant nondegenerate bilinear form  $\langle \cdot, \cdot \rangle$ . A Frobenius manifold is a manifold  $M$  with a smooth structure of a Frobenius algebra on the tangent space  $T_t M$  at any point  $t \in M$  with certain compatibility conditions [6]. Globally, we require the metric  $\langle \cdot, \cdot \rangle$  to be flat and the unity vector field  $e$  to be covariantly constant with respect to it. In the flat coordinates  $(t^1, \dots, t^r)$  where

$e = \frac{\partial}{\partial t^r}$  the compatibility conditions imply that there exists a function  $F(t^1, \dots, t^r)$  such that

$$\eta_{ij} = \langle \partial_i, \partial_j \rangle = \partial_r \partial_i \partial_j F(t)$$

and the structure constants of the Frobenius algebra are given by

$$C^k_{ij} = \sum_p \Omega_1^{kp} \partial_{t^p} \partial_{t^i} \partial_{t^j} F(t)$$

where  $\Omega_1^{ij}$  denotes the inverse of the matrix  $\eta_{ij}$ . In this work, we consider Frobenius manifolds where the quasihomogeneity condition takes the form

$$\sum_{i=1}^r d_i t^i \partial_{t^i} F(t) = (3-d)F(t); d_r = 1. \tag{2.1}$$

This condition defines the degrees  $d_i$  and the charge  $d$  of the Frobenius structure. The associativity of the Frobenius algebra implies that the potential  $F(t)$  satisfies a system of partial differential equations which appears in topological field theory and is called Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) equations:

$$\sum_{k,p} \partial_{t^i} \partial_{t^j} \partial_{t^k} F(t) \Omega_1^{kp} \partial_{t^p} \partial_{t^q} \partial_{t^n} F(t) = \sum_{k,p} \partial_{t^n} \partial_{t^j} \partial_{t^k} F(t) \Omega_1^{kp} \partial_{t^p} \partial_{t^q} \partial_{t^i} F(t), \forall i, j, q, n. \tag{2.2}$$

Detailed information about Frobenius manifolds and related topics can be found in [1].

### 2.2 Flat pencil of metrics and Frobenius manifolds

In this section we review the relation between the geometry of flat pencil of metrics and Frobenius manifolds. See [15] for details.

Let  $M$  be a smooth manifold of dimension  $r$ . A symmetric bilinear form  $(\cdot, \cdot)$  on  $T^*M$  is called a contravariant metric if it is invertible on an open dense subset  $M_0 \subset M$ . In local coordinates, if we set

$$\Omega^{ij}(u) = (du^i, du^j); i, j = 1, \dots, r \quad (2.3)$$

then the inverse matrix  $\Omega_{ij}(u)$  of  $\Omega^{ij}(u)$  determines a metric  $\langle \cdot, \cdot \rangle$  on  $TM_0$ . We define the contravariant Christoffel symbols  $\Gamma_k^{ij}$  of  $(\cdot, \cdot)$  by  $\Gamma_k^{ij} := -\sum_s \Omega^{is} \Gamma_{sk}^j$  where  $\Gamma_{sk}^j$  are the Christoffel symbols of  $\langle \cdot, \cdot \rangle$ . We say the metric  $(\cdot, \cdot)$  is flat if  $\langle \cdot, \cdot \rangle$  is flat.

Let  $(\cdot, \cdot)_1$  and  $(\cdot, \cdot)_2$  be two contravariant flat metrics on  $M$  and denote their Christoffel symbols by  $\Gamma_{1;k}^{ij}(u)$  and  $\Gamma_{2;k}^{ij}(u)$  respectively. We say  $(\cdot, \cdot)_1$  and  $(\cdot, \cdot)_2$  form a flat pencil of metrics if  $(\cdot, \cdot)_\lambda := (\cdot, \cdot)_1 + \lambda(\cdot, \cdot)_2$  defines a flat metric on  $T^*M$  for a generic  $\lambda$  and its Christoffel symbols are given by  $\Gamma_{\lambda;k}^{ij}(u) = \Gamma_{2;k}^{ij}(u) + \Gamma_{1;k}^{ij}(u)$ .

Let  $(\cdot, \cdot)_1$  and  $(\cdot, \cdot)_2$  be two contravariant metrics on  $M$  and denote their matrices by  $\Omega_1^{ij}(u)$  and  $\Omega_2^{ij}(u)$ , respectively, in some coordinates  $(u^1, \dots, u^r)$ . Suppose that they form a flat pencil of metrics. This flat pencil of metrics is called *quasihomogeneous of degree  $d$*  if there exists a function  $\square$  on  $M$  such that the vector fields

$$\begin{aligned} E &:= \nabla_2 \tau, & E^i &= \sum_s \Omega_2^{is} \partial_s \tau \\ e &:= \nabla_1 \tau, & e^i &= \sum_s \Omega_1^{is} \partial_s \tau \end{aligned} \quad (2.4)$$

satisfy the following relations

$$[e, E] = e, \text{Lie}_E(\cdot)_2 = (d-1)(\cdot)_2, \text{Lie}_e(\cdot)_2 = (\cdot)_1, \text{Lie}_e(\cdot)_1 = 0.$$

Here  $\text{Lie}_X$  denote the Lie derivative along a given vector field  $X$ . In addition, the quasihomogeneous flat pencil of metrics is called *regular* if the (1,1)-tensor  $R_{ij}^j = \frac{d-1}{2} \delta_i^j + \nabla_{1i} E^j$  is nondegenerate on  $M$ .

The following theorem due to Dubrovin gives a connection between the geometry of Frobenius manifolds and flat pencils of metrics.

**Theorem 2.1.** [15] *A quasihomogeneous regular flat pencil of metrics of degree  $d$  on a manifold  $M$  defines a Frobenius structure on  $M$  of charge  $d$ .*

Let us assume the flat pencil of metrics on  $M$  is regular quasihomogeneous of degree  $d$ . Let  $(t^1, \dots, t^r)$  be flat coordinates of  $(\cdot, \cdot)_1$  where  $\tau = t^1, e = \partial_{t^r}$  and  $E = \sum_i d_i t^i \partial_{t^i}$ . Let  $\eta_{ij}$  denote the inverse of  $\Omega_1^{ij}(t)$ . Then it turns

out that the potential  $\mathbb{F}(t^1, \dots, t^r)$  is obtained from the equations

$$\begin{aligned} \frac{\partial^2 \mathbb{F}}{\partial t^i \partial t^j} &= \sum_{k,l} \frac{1}{d-1+d_k+d_l} \eta_{ik} \eta_{jl} \Omega_2^{kl}(t) \\ \frac{\partial \mathbb{F}}{\partial t^i} &= \frac{1}{(3-d-d_i)} \sum_k d_k t^k \frac{\partial^2 \mathbb{F}}{\partial t^k \partial t^i} \\ \mathbb{F}(t) &= \frac{1}{3-d} \sum_k d_k t^k \frac{\partial \mathbb{F}}{\partial t^k} \end{aligned} \quad (2.5)$$

It is well known that from a Frobenius manifold we always have a flat pencil of metrics but it does not necessarily satisfy the regularity condition [15].

### 3. Dicyclic groups

Let  $n$  be a natural number greater than 1 and  $W$  be the matrix group generated by

$$\sigma := \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix}, \alpha := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (3.1)$$

where  $\xi$  is a primitive  $2n$ -th root of unity. Then  $\sigma$  and  $\alpha$  satisfy the relations

$$\sigma^{2n} = 1, \alpha^2 = -\sigma^n, \alpha^{-1} \sigma \alpha = \sigma^{-1}. \quad (3.2)$$

Thus  $W$  is isomorphic to the dicyclic group of order  $4n$ . The invariant ring of  $W$  is generated by the following homogeneous polynomials [2]

$$u_1 = x_1^2 x_2^2, u_2 = x_1^{2n} + x_2^{2n}, u_3 = x_1 x_2 (x_1^{2n} - x_2^{2n}) \quad (3.3)$$

subject to the relation

$$u_3^2 - u_1 u_2^2 + 4u_1^{n+1} = 0. \quad (3.4)$$

The orbits space  $M$  of  $W$  is a variety isomorphic to the hypersurface  $T$  defined as the zero set of equation (3.4) in  $\mathbb{C}^3$ . Consider equation (3.4) as a quadratic equation in  $u_3$ . Then any point  $p$  out of the discriminant locus has a small

neighbourhood  $U_p$  where  $u_1$  and  $u_2$  act as coordinates. In what follows we assume that we fix such the open set  $U \subset V$  with coordinates  $(u_1, u_2)$  and we omit the subscript  $p$ .

Let  $h$  be the Hessian matrix of  $u_1$ , i.e.  $h_{ij} = \frac{\partial^2 u_1}{\partial x_i \partial x_j}$  and let  $h^{-1}$  denotes its inverse. Then, by direct calculations,  $h^{-1}$  defines a flat contravariant metric  $(\cdot, \cdot)_2$  on  $U$ . This metric, in the coordinates  $u_1$  and  $u_2$ , is given as follows

$$(\cdot, \cdot)_2 = \begin{pmatrix} \frac{4}{3}u_1 & \frac{2n}{3}u_2 \\ \frac{2n}{3}u_2 & -\frac{2n^2}{3u_1}(u_2^2 - 6u_1^n) \end{pmatrix}; \quad (dt_i, dt_j)_2 = \sum_{k,l=1}^2 \frac{\partial u_i}{\partial x_k} \frac{\partial u_j}{\partial x_l} h_{kl}^{-1}. \quad (3.5)$$

Let  $e$  be a vector field of the form  $f(u_1)\partial_{u_2}$ , where  $f(u_1)$  is any smooth function. Then, by direct calculations, the Lie derivative  $(\cdot, \cdot)_1$  of  $(\cdot, \cdot)_2$  along  $e$  forms with  $(\cdot, \cdot)_2$  a flat pencil of metrics. This metric takes the value

$$(\cdot, \cdot)_1 = \begin{pmatrix} 0 & \frac{2}{3}(nf - 2u_1 f') \\ \frac{2}{3}(nf - 2u_1 f') & -\frac{4}{3u_1}(n^2 u_2 f + nu_1 u_2 f') \end{pmatrix}. \quad (3.6)$$

The guess for the vector field to take this from was inspired by [1]. In order to get a quasihomogeneous flat pencil of metrics, we need the Lie derivative of  $(\cdot, \cdot)_1$  with respect to  $e$  to equal zero. This condition leads to the following differential equation for  $f(u)$

$$2nu_1 f f' - 2u_1^2 (f')^2 + n^2 f^2 = 0 \quad (3.7)$$

which has two independent solutions

$$f_+ = u_1^{\frac{n}{2}(1+\sqrt{3})} \quad \text{and} \quad f_- = u_1^{\frac{n}{2}(1-\sqrt{3})}. \quad (3.8)$$

Let us assume  $e = f_+ \partial_{u_2} = u_1^{\frac{n}{2}(1+\sqrt{3})} \partial_{u_2}$ . Then

$$(dt_i, dt_j)_1 = \begin{pmatrix} 0 & -\frac{2n}{\sqrt{3}} u_1^{\frac{1}{2}(1+\sqrt{3})n} \\ -\frac{2n}{\sqrt{3}} u_1^{\frac{1}{2}(1+\sqrt{3})n} & -\frac{2}{3}(3+\sqrt{3})n^2 u_1^{\frac{1}{2}(1+\sqrt{3})n-1} u_2 \end{pmatrix}. \quad (3.9)$$

It turns out that the two metrics  $(\cdot, \cdot)_2$  and  $(\cdot, \cdot)_1$  form a quasihomogeneous flat pencil of metrics with degree

$$d = \frac{2 + \sqrt{3}n}{\sqrt{3}n}. \quad (3.10)$$

In the notations of equations (2.4), we have  $\tau = -\frac{\sqrt{3}}{2n}u_1$  and

$$E = -\frac{2}{\sqrt{3}n}u_1 \partial_{u_1} - \frac{1}{\sqrt{3}}u_2 \partial_{u_2}. \quad (3.11)$$

This flat pencil of metrics is also regular since the  $(1,1)$ -tensor  $R^i_j$  equals the nondegenerate matrix

$$\begin{pmatrix} -\frac{1}{\sqrt{3}n} & 0 \\ 0 & \frac{1-n}{\sqrt{3}n} \end{pmatrix}. \quad (3.12)$$

Flat coordinates for  $(\cdot, \cdot)_1$  are obtained by setting

$$t_1 = -\frac{\sqrt{3}}{2n}u_1, \quad t_2 = u_2 u_1^{-\frac{1}{2}(1+\sqrt{3})n}. \quad (3.13)$$

In these coordinates we get

$$(\cdot, \cdot)_2 = \begin{pmatrix} -\frac{2}{\sqrt{3}n}t_1 & t_2 \\ t_2 & 2^{1-\sqrt{3}n} 3^{\frac{1}{2}(\sqrt{3}n+1)} n^2 (-nt_1)^{-\sqrt{3}n-1} \end{pmatrix}, \quad (\cdot, \cdot)_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (3.14)$$

The potential  $\mathbb{F}_+$  of the corresponding Frobenius manifold is

$$\mathbb{F}_+ = \frac{2^{-\sqrt{3}n} 3^{\frac{1}{2}(\sqrt{3}n+1)} (-nt_1)^{1-\sqrt{3}n}}{3n^2 - 1} + \frac{1}{2}t_1 t_2^2. \quad (3.15)$$

Let us take  $e = f_- \partial_{u_2} = u_1^{\frac{n}{2}(1-\sqrt{3})} \partial_{u_2}$ . Then similar to the method above, we get a regular quasihomogenous flat pencil of metrics of degree

$$d = \frac{2 - \sqrt{3}n}{\sqrt{3}n} \quad (3.16)$$

with  $\tau = \frac{\sqrt{3}}{2n}u_1$ . The resulting potential will be

$$\mathbb{F}_- = \frac{2^{\sqrt{3}n} 3^{\frac{1}{2}(\sqrt{3}n-1)} (nt_1)^{\sqrt{3}n+1}}{3n^2 - 1} + \frac{1}{2}t_1 t_2^2. \quad (3.17)$$

We repeat the calculation by taking  $(u_1, u_3)$  as coordinates instead of  $(u_1, u_2)$ . It turns out that even though the middle steps may differ in values, the resulting Frobenius manifolds are exactly the same as those given by the potentials (3.15),(3.17).

We observe that Dubrovin computed by an ad hoc procedure all possible potentials of 2-dimensional Frobenius manifolds [6]. The potentials found in this article, after scaling, are listed by Dubrovin in the form

$$\mathbb{F}(z_1, z_2) = z_1^k + \frac{1}{2}z_2^2z_1, \quad k = \frac{3-d}{1-d} \quad (3.18)$$

where  $d$  is  $\frac{2+\sqrt{3n}}{\sqrt{3n}}$  or  $\frac{2-\sqrt{3n}}{\sqrt{3n}}$ . However, finding this by using the method of a flat pencil of metrics on an orbits space of a finite group that is not a reflection group is a surprising result.

#### 4. Conclusion

In this paper we prove that for a linear representation of Dicyclic groups we can still use Dubrovin's method to construct Frobenius manifolds on its orbits spaces. In addition, the method leads to two structures of Frobenius manifolds. The result reported in this article is a part of work in progress to apply Dubrovin's method on orbits spaces of finite groups to find interesting new examples of Frobenius manifolds. In future, we will consider irreducible representations of Coxeter groups which are not reflection representations [16].

#### Conflict of interest

The authors declare no conflict of interest.

#### Acknowledgment

This work is funded by the internal grant of Sultan Qaboos University (IG/SCI/DOMS/19/08). The authors thank Hans-Christian Herbig for stimulating discussions and the anonymous reviewers for their comments and suggestions.

#### References

1. Dubrovin, B. Geometry of 2D topological field theories. Integrable systems and quantum groups, *Montecatini Terme*, 1993, Lecture Notes in Math., Springer, Berlin, 1996, **1620**, 120-348.
2. Neusel, M. *Invariant Theory*, American Mathematical Society, Student Mathematical Library (Book 36), ISBN: 978-0821841327, 2006,
3. Derksen, H. and Kemper, G. *Computational invariant theory. Invariant Theory and Algebraic Transformation Groups*, I. Encyclopaedia of Mathematical Sciences, **130**. Springer-Verlag, Berlin, 2002.
4. Orlik, P. and Solomon, L. The hessian map in the invariant theory of reflection groups, *Nagoya Mathematical Journal*, 1988, **109**, 1-21.
5. Saito, K. On a linear structure of the quotient variety by a finite reflexion group. *Publications of the Research Institute for Mathematical Sciences*, 1993, **29(4)**, 535-579.
6. Saito, K., Yano, T. and Sekeguchi, J. On a certain generator system of the ring of invariants of a finite reflection group, *Communication in Algebra*, 1980, **8(4)**, 373-408.
7. Dubrovin, B. On almost duality for Frobenius manifolds, *Geometry, topology and mathematical physics*, American Mathematical Society, 2004, Ser.2, vol.**212**, 75-132.
8. Dubrovin, B. Differential geometry of the space of orbits of a Coxeter group, *Surveys in differential geometry IV: Integrable systems*, International Press, 1998, 181-211.
9. Zuo, D. Frobenius manifolds associated to  $B_1$  and  $D_1$ , revisited, *International Mathematics Research Notices*, 2007, **2007**, rnm020.
10. Dubrovin, B. and Zhang, Y. Extended affine Weyl groups and Frobenius manifolds, *Compositio Mathematica*, 1998, **111**, 167-219.
11. Dubrovin, B., Strachan, Ian A.B., Zhang, Y. and Zuo, D. Extended affine Weyl groups of BCD-type: their Frobenius manifolds and Landau-Ginzburg superpotentials, *Advances in Mathematics*, 2019, 351.
12. Bertola, M. Jacobi groups, Jacobi forms and their applications. PhD. Thesis, SISSA, Trieste 1999.
13. Zuo, D. Frobenius manifolds and a new class of extended affine Weyl groups  $W^{(k,k+1)}(A_1)$ . *Letters in Mathematical Physics*, 2020, **110 (7)**, 1903-1940.
14. Arsie, A. and Lorenzoni, P. Complex reflection groups, logarithmic connections and bi-flat F-manifolds. *Letters in Mathematical Physics*, 2017, **107(10)**, 1919-1961.
15. Dubrovin, B. Flat pencils of metrics and Frobenius manifolds. *Integrable systems and algebraic geometry Kobe/Kyoto*, 1997, 47-72.
16. Al-Maamari, Z. and Dinar, Y. Frobenius manifolds from a few linear representations, in preparation.

---

Received 15 November 2019

Accepted 16 July 2020